


## Angola


General indicators	Source	2001
Population (thousands)	UNPOP	13527
Population growth rate (%)	UNPOP	3.2
Infant mortality rate (per 1000)	UNICEF	154
Under-5 mortality rate (per 1000)	UNICEF	260
Crude birth rate (per 1000)	UNPOP	51.3
Pop. at endemic (epidemic) risk (%)	MARA <sup>1</sup>	90 (8)

### Proportional rank for select indicators



### Malaria morbidity and mortality


Hospital deaths due to malaria	25
Malaria admissions	20
Malaria outpatient attendance	50
Outpatient malaria cases (per 1000)	94
Inpatient malaria cases (per 1000)	62



Source	1998	1999	2000	2001
MoH	25			
MoH		20		
MoH		50		
MoH	94	62	105	66

### Mosquito nets and ITNs


Households with mosquito nets	
Under-5s using mosquito nets	10
Households with ITNs	
Under-5s using ITNs	2



Source	1998	1999	2000	2001
MICS				10
MICS				2

### Antimalarial treatment


Under-5s receiving any antimalarial	63
Under-5s receiving chloroquine	57
Under-5s receiving any antimalarial <24 h	



Source	1998	1999	2000	2001
MICS				63
MICS				57

### Attending health facility


Under-5s receiving any antimalarial	45
Under-5s receiving chloroquine	40




Source	1998	1999	2000	2001
MICS				45
MICS				40

### Malaria in pregnancy

PW receiving ANC at least once	
PW attending ANC at least twice	
PW receiving IPT at least once	
PW receiving IPT at least twice	
PW using mosquito nets	
PW using ITNs	



Source	1998	1999	2000	2001
MICS				66



### Antimalarial treatment policy

Date adopted:	----	
Uncomplicated	Probable	CQ 25 mg/kg
	Confirmed	CQ 25 mg/kg
Treatment failure		SP
Severe malaria		Quinine 3 times a day, 7 days
Pregnancy	Prevention	CQ weekly

### Taxes and tariffs on insecticide-treated nets

	Date
Import tariff - nets	
VAT - nets	

### Notes

For an explanation of indicators, ranks, and other information, see the Country Profile Overview.

Sources listed reflect most recent value.

Case notification rate is based on number of reported outpatient cases.

Nationally representative surveys (MICS and DHS) are used for reporting intervention coverage estimates for 1) mosquito net and ITN possession and use and 2) antimalarial treatment.

<sup>1</sup> Population at risk is determined from modelling retrospective climate data and population projections. Percentage of population at risk does not vary by year.