


Burkina Faso

General indicators	Source	2001
Population (thousands)	UNPOP	11856
Population growth rate (%)	UNPOP	2.5
Infant mortality rate (per 1000)	UNICEF	104
Under-5 mortality rate (per 1000)	UNICEF	197
Crude birth rate (per 1000)	UNPOP	46.8
Pop. at endemic (epidemic) risk (%)	MARA ¹	100 (0)

Proportional rank for select indicators



Malaria morbidity and mortality

Hospital deaths due to malaria	30%
Malaria admissions	25%
Malaria outpatient attendance	30%
Outpatient malaria cases (per 1000)	81
Inpatient malaria cases (per 1000)	3

0% 20% 40% 60% 80% 100%

Source	1998	1999	2000	2001
MoH	33	31		
MoH	26	27		
MoH	30	31		
MoH	81	78	6	
MoH	3	3		

Mosquito nets and ITNs

Households with mosquito nets	30%
Under-5s using mosquito nets	25%
Households with ITNs	30%
Under-5s using ITNs	25%

0% 20% 40% 60% 80% 100%

Source	1998	1999	2000	2001
--------	------	------	------	------

Antimalarial treatment

Under-5s receiving any antimalarial	30%
Under-5s receiving chloroquine	25%
Under-5s receiving any antimalarial <24 h	30%

0% 20% 40% 60% 80% 100%

Source	1998	1999	2000	2001
--------	------	------	------	------

Attending health facility


Under-5s receiving any antimalarial	30%
Under-5s receiving chloroquine	25%

0% 20% 40% 60% 80% 100%

Malaria in pregnancy

PW receiving ANC at least once	60%
PW attending ANC at least twice	50%
PW receiving IPT at least once	30%
PW receiving IPT at least twice	25%
PW using mosquito nets	30%
PW using ITNs	25%

Source	1998	1999	2000	2001
DHS		63		
DHS		56		



Antimalarial treatment policy

Date adopted: ----

Uncomplicated	Probable	CQ 25 mg/kg
	Confirmed	CQ 25 mg/kg
Treatment failure		SP
Severe malaria		Quinine 3 times a day, 7 days
Pregnancy	Prevention	CQ weekly

Taxes and tariffs on insecticide-treated nets

		Date
Import tariff - nets	18%	2000
VAT - nets	18%	2000

Notes

For an explanation of indicators, ranks, and other information, see the Country Profile Overview.

Sources listed reflect most recent value.

Case notification rate is based on number of reported outpatient cases.

Nationally representative surveys (MICS and DHS) are used for reporting intervention coverage estimates for 1) mosquito net and ITN possession and use and 2) antimalarial treatment.

¹ Population at risk is determined from modelling retrospective climate data and population projections. Percentage of population at risk does not vary by year.