


Guinea

General indicators	Source	2001
Population (thousands)	UNPOP	8274
Population growth rate (%)	UNPOP	2.7
Infant mortality rate (per 1000)	UNICEF	109
Under-5 mortality rate (per 1000)	UNICEF	169
Crude birth rate (per 1000)	UNPOP	43.4
Pop. at endemic (epidemic) risk (%)	MARA ¹	100 (<1)

Proportional rank for select indicators



Malaria morbidity and mortality

Hospital deaths due to malaria	15%
Malaria admissions	25%
Malaria outpatient attendance	35%
Outpatient malaria cases (per 1000)	104
Inpatient malaria cases (per 1000)	2

0% 20% 40% 60% 80% 100%

Source	1998	1999	2000	2001
MoH	1	1	26	15
MoH	23	23	23	
MoH	27	27	33	30

Mosquito nets and ITNs

Households with mosquito nets	0%
Under-5s using mosquito nets	0%
Households with ITNs	0%
Under-5s using ITNs	0%

0% 20% 40% 60% 80% 100%

Source	1998	1999	2000	2001
--------	------	------	------	------

Antimalarial treatment

Under-5s receiving any antimalarial	0%
Under-5s receiving chloroquine	0%
Under-5s receiving any antimalarial <24 h	0%

0% 20% 40% 60% 80% 100%

Source	1998	1999	2000	2001
--------	------	------	------	------

Attending health facility


Under-5s receiving any antimalarial	0%
Under-5s receiving chloroquine	0%

Malaria in pregnancy

PW receiving ANC at least once	85%
PW attending ANC at least twice	75%
PW receiving IPT at least once	0%
PW receiving IPT at least twice	0%
PW using mosquito nets	0%
PW using ITNs	0%

0% 20% 40% 60% 80% 100%

Source	1998	1999	2000	2001
DHS		74		
DHS		69		



Antimalarial treatment policy

Date adopted:	----	
Uncomplicated	Probable	CQ 25 mg/kg
	Confirmed	CQ 25 mg/kg
Treatment failure		SP
Severe malaria		Quinine 3 times a day, 7 days
Pregnancy	Prevention	CQ weekly

Taxes and tariffs on insecticide-treated nets

		Date
Import tariff - nets	21%	2000
VAT - nets	21%	2000

Notes

For an explanation of indicators, ranks, and other information, see the Country Profile Overview.

Sources listed reflect most recent value.

Case notification rate is based on number of reported outpatient cases.

Nationally representative surveys (MICS and DHS) are used for reporting intervention coverage estimates for 1) mosquito net and ITN possession and use and 2) antimalarial treatment.

¹ Population at risk is determined from modelling retrospective climate data and population projections. Percentage of population at risk does not vary by year.