


## Mozambique

General indicators	Source	2001
Population (thousands)	UNPOP	18644
Population growth rate (%)	UNPOP	2.9
Infant mortality rate (per 1000)	UNICEF	125
Under-5 mortality rate (per 1000)	UNICEF	197
Crude birth rate (per 1000)	UNPOP	41.7
Pop. at endemic (epidemic) risk (%)	MARA <sup>1</sup>	100 (<1)

### Proportional rank for select indicators



### Malaria morbidity and mortality

Hospital deaths due to malaria	20%
Malaria admissions	40%
Malaria outpatient attendance	40%
Outpatient malaria cases (per 1000)	
Inpatient malaria cases (per 1000)	

0% 20% 40% 60% 80% 100%

Source	1998	1999	2000	2001
MoH	24			
MoH	28	48	44	
MoH	11	130	179	198

### Mosquito nets and ITNs

Households with mosquito nets	
Under-5s using mosquito nets	
Households with ITNs	
Under-5s using ITNs	

0% 20% 40% 60% 80% 100%

Source	1998	1999	2000	2001
--------	------	------	------	------

### Antimalarial treatment

Under-5s receiving any antimalarial	
Under-5s receiving chloroquine	
Under-5s receiving any antimalarial <24 h	

0% 20% 40% 60% 80% 100%

Source	1998	1999	2000	2001
--------	------	------	------	------

### Attending health facility

Under-5s receiving any antimalarial	
Under-5s receiving chloroquine	

0% 20% 40% 60% 80% 100%


Source	1998	1999	2000	2001
--------	------	------	------	------

### Malaria in pregnancy

PW receiving ANC at least once	
PW attending ANC at least twice	
PW receiving IPT at least once	
PW receiving IPT at least twice	
PW using mosquito nets	
PW using ITNs	

0% 20% 40% 60% 80% 100%

Source	1998	1999	2000	2001
--------	------	------	------	------



### Antimalarial treatment policy

Date adopted:	2002	
Uncomplicated	Probable	CQ 25 mg/kg
	Confirmed	CQ 25 mg/kg
Treatment failure	SP	
Severe malaria	Quinine 3 times a day, 7 days	
Pregnancy	Prevention	CQ weekly

### Taxes and tariffs on insecticide-treated nets

	Date
Import tariff - nets	
VAT - nets	

### Notes

For an explanation of indicators, ranks, and other information, see the Country Profile Overview.

Sources listed reflect most recent value.

Case notification rate is based on number of reported outpatient cases.

Nationally representative surveys (MICS and DHS) are used for reporting intervention coverage estimates for 1) mosquito net and ITN possession and use and 2) antimalarial treatment.

<sup>1</sup> Population at risk is determined from modelling retrospective climate data and population projections. Percentage of population at risk does not vary by year.