


Togo

General indicators	Source	2001
Population (thousands)	UNPOP	4657
Population growth rate (%)	UNPOP	2.8
Infant mortality rate (per 1000)	UNICEF	79
Under-5 mortality rate (per 1000)	UNICEF	141
Crude birth rate (per 1000)	UNPOP	38.7
Pop. at endemic (epidemic) risk (%)	MARA ¹	100 (0)

Proportional rank for select indicators



Malaria morbidity and mortality

Hospital deaths due to malaria	20%
Malaria admissions	25%
Malaria outpatient attendance	40%
Outpatient malaria cases (per 1000)	87
Inpatient malaria cases (per 1000)	2

0% 20% 40% 60% 80% 100%

Source	1998	1999	2000	2001
MoH	23	27	19	20
MoH	26	24	19	22
MoH	39	38	39	39
MoH	87	94	88	93
MoH	2	3	3	3

Mosquito nets and ITNs

Households with mosquito nets	20%
Under-5s using mosquito nets	15%
Households with ITNs	2%
Under-5s using ITNs	2%

0% 20% 40% 60% 80% 100%

Source	1998	1999	2000	2001
MICS			15	
MICS			2	

Antimalarial treatment

Under-5s receiving any antimalarial	60%
Under-5s receiving chloroquine	59%
Under-5s receiving any antimalarial <24 h	

0% 20% 40% 60% 80% 100%

Source	1998	1999	2000	2001
MICS			60	
MICS			59	

Attending health facility

Under-5s receiving any antimalarial	21%
Under-5s receiving chloroquine	21%

0% 20% 40% 60% 80% 100%


Source	1998	1999	2000	2001
MICS			21	
MICS			21	

Malaria in pregnancy

PW receiving ANC at least once	73%
PW attending ANC at least twice	
PW receiving IPT at least once	
PW receiving IPT at least twice	
PW using mosquito nets	
PW using ITNs	

0% 20% 40% 60% 80% 100%

Source	1998	1999	2000	2001
MICS			73	



Antimalarial treatment policy

Date adopted:	----	
Uncomplicated	Probable	CQ 25 mg/kg
	Confirmed	CQ 25 mg/kg
Treatment failure		SP
Severe malaria		Quinine 3 times a day, 7 days
Pregnancy	Prevention	CQ weekly

Taxes and tariffs on insecticide-treated nets

		Date
Import tariff - nets	18%	2000
VAT - nets	18%	2000

Notes

For an explanation of indicators, ranks, and other information, see the Country Profile Overview.

Sources listed reflect most recent value.

Case notification rate is based on number of reported outpatient cases.

Nationally representative surveys (MICS and DHS) are used for reporting intervention coverage estimates for 1) mosquito net and ITN possession and use and 2) antimalarial treatment.

¹ Population at risk is determined from modelling retrospective climate data and population projections. Percentage of population at risk does not vary by year.