


Zimbabwe

General indicators	Source	2001
Population (thousands)	UNPOP	12852
Population growth rate (%)	UNPOP	2.0
Infant mortality rate (per 1000)	UNICEF	76
Under-5 mortality rate (per 1000)	UNICEF	123
Crude birth rate (per 1000)	UNPOP	35.1
Pop. at risk (%)	MARA ¹	84

Proportional rank for select indicators



Malaria morbidity and mortality

Hospital deaths due to malaria	10%
Malaria admissions	15%
Malaria outpatient attendance	15%
Outpatient malaria cases (per 1000)	141
Inpatient malaria cases (per 1000)	96

0% 20% 40% 60% 80% 100%

Source	1998	1999	2000	2001
MoH	8			
MoH	15		20	
MoH	12		15	
MoH	141	96	121	
MoH			1	

Mosquito nets and ITNs

Households with mosquito nets	10%
Under-5s using mosquito nets	3%
Households with ITNs	
Under-5s using ITNs	

0% 20% 40% 60% 80% 100%

Source	1998	1999	2000	2001
DHS		10		
DHS		3		

Antimalarial treatment

Under-5s receiving any antimalarial	
Under-5s receiving chloroquine	
Under-5s receiving any antimalarial <24 h	

0% 20% 40% 60% 80% 100%

Source	1998	1999	2000	2001
--------	------	------	------	------

Attending health facility


Under-5s receiving any antimalarial	
Under-5s receiving chloroquine	

Malaria in pregnancy

PW receiving ANC at least once	93%
PW attending ANC at least twice	80%
PW receiving IPT at least once	
PW receiving IPT at least twice	
PW using mosquito nets	
PW using ITNs	

0% 20% 40% 60% 80% 100%

Source	1998	1999	2000	2001
DHS		93		
DHS		80		



Antimalarial treatment policy

	Date adopted:	2001
Uncomplicated	Probable	CQ + SP
	Confirmed	CQ + SP
Treatment failure		Quinine 3 times a day, 7 days
Severe malaria		Quinine 3 times a day, 7 days
Pregnancy	Prevention	Maloprim weekly

Taxes and tariffs on insecticide-treated nets

		Date
Import tariff - nets	10%	Before Abuja
VAT - nets	20%	Before Abuja

Notes

For an explanation of indicators, ranks, and other information, see the Country Profile Overview. Sources listed reflect most recent value.

Case notification rate is based on number of reported outpatient cases.

Nationally representative surveys (MICS and DHS) are used for reporting intervention coverage estimates for 1) mosquito net and ITN possession and use and 2) antimalarial treatment.

¹ Population at risk is determined from modelling retrospective climate data and population projections. Percentage of population at risk does not vary by year.