

Table A.12 Percentage of pregnant women that slept under an insecticide-treated net during the night preceding the survey, by background characteristics

Africa	Year	Source	Scale	Total	Residence		Wealth quintile					
					Urban	Rural	Poorest	Second	Middle	Fourth	Least poor	
Benin	2004	MoH/RBM 2004	6 health zones	28.7	-	-	-	-	-	-	-	
	2001	RBM 2000	3 health zones	3.8	-	-	-	-	-	-	-	
Burkina Faso	2003	DHS 2003	national	2.6	6.1	2.1	0.6	0.7	2.7	1.5	8.3	
	2001	RBM 2000	district	10.0	-	-	-	-	-	-	-	
Chad	2000	RBM 2001	5 districts	7.2	-	-	-	-	-	-	-	
Eritrea	2002	DHS 2002	national	2.9	-	-	3.4	2.3	0.8	3.8	5.5	
Ghana	2003	DHS 2003	national	2.7	1.6	3.2	4.7	2.8	1.0	1.9	3.0	
	2001	RBM 2001	5 districts	7.8	-	-	-	-	-	-	-	
Guinea	2001	RBM 2001	4 districts	2.7	-	-	-	-	-	-	-	
Kenya	2003	DHS 2003	national	4.4	4.8	4.3	1.7	2.2	6.5	6.3	5.9	
Malawi	2004	MoH 2004	national	31.4	49.1	29.0	-	-	-	-	-	
Mali	2003	NetMark 2003	5 areas	19.6	-	-	-	-	-	-	-	
		RBM 2003	district	19.0	-	-	-	-	-	-	-	
Mozambique	2000	NetMark 2000	5 areas	5.6	-	-	-	-	-	-	-	
Nigeria	2003	DHS 2003	national	1.3	0.4	1.6	-	-	-	-	-	
	2000	NetMark 2000	5 areas	0.0	-	-	-	-	-	-	-	
Senegal	2000	NetMark 2000	5 areas	6.0	-	-	-	-	-	-	-	
Uganda	2001	DHS 2000–2001	national	0.5	-	-	-	-	-	-	-	
		MoH 2001 (RBM baseline survey)	4 districts	2.3	-	-	-	-	-	-	-	
	2000	NetMark 2000	5 areas	1.5	-	-	-	-	-	-	-	
Zambia	2002	DHS 2002–2003	national	7.9	-	-	-	-	-	-	-	
	2000	NetMark 2000	5 areas	1.4	-	-	-	-	-	-	-	
The Americas												
	Year	Source	Scale	Total								
Bolivia	2001	PSI 2001a	1 province	17.8								