

Table A.17 Pregnant women receiving SP at least once during pregnancy (community level, prevention or treatment), by background characteristics

Africa	Year	Source	Scale	Total	Residence		Wealth quintile				
					Urban	Rural	Poorest	Second	Middle	Fourth	Least poor
Benin	2001	DHS 2001	national	6.2	4.2	7.2	-	-	-	-	-
Ghana	2003	DHS 2003	national	1.0	0.9	1.1	1.3	0.7	1.0	0.9	1.4
Kenya	2003	DHS 2003	national	12.5	13.1	12.4	12.6	11.8	13.8	11.9	12.5
Malawi	2004	MoH 2004	national	77.7	88.9	75.4	-	-	-	-	-
	2000	DHS 2000	national	67.5	-	-	-	-	-	-	-
Mauritania	2004	DHS 2003–2004	national	0.5	0.9	0.2	-	-	-	-	-
Nigeria	2003	DHS 2003	national	2.4	-	-	-	-	-	-	-
Rwanda	2000	DHS 2000	national	0.1	-	-	-	-	-	-	-
Zambia	2002	DHS 2002–2003	national	0.5	0.8	0.4	-	-	-	-	-

Table A.18 Pregnant women receiving SP at least twice during pregnancy (community level, prevention or treatment), by background characteristics

Africa	Year	Source	Scale	Total	Residence	
					Urban	Rural
Malawi	2004	MoH 2004	national	46.8	57.2	44.7
	2000	DHS 2000	national	29.3	-	-